



WHAT IS SPRYCEL® DASATINIB ?

- Cancer drug used in the treatment of some types of blood cancers (leukaemias)
- Prescribed and renewed by your hospital doctor specialized in oncology, cancerology or hematology.
- Available in hospital pharmacy on presentation of your prescription.




- Oral suspension in white to opaque yellow colour, dosed at **10 mg/ml**.
- *The reconstitution of the oral suspension is done by a pharmacist or a qualified health care professional, before dispensing.*
- Bottle provided with a syringe for 12 ml oral dispensing.



- Store it in the fridge (between 2° and 8°C) for 60 days maximum
- Keep the bottle upright, out of sight and reach of children.
- Wash your hands carefully before and after each handling of the suspension.
- Do not throw away the opened bottles, please bring them back to your pharmacist.

HOW TO TAKE THIS MEDICINE?



Dose per day mg = mL
When ?h..... = 1 take a day, during or away from meals. <i>May be mixed with milk, yogourt, apple juice or compote</i>
How ?	 <p>① Shake the bottle (approx. 30"), take off the cap and check that the adapter is well set into the bottle. ② See that the 12ml syringe piston supplied is thrust to its maximum, then insert the syringe tip into the adapter while holding it vertically. ③ Keep the tip of the syringe well set into the bottle, then turn it upside down with the syringe turned downward. ④ Draw the number of ml necessary by slowly pulling the piston. ⑤ Keep the syringe tip well thrust into the bottle and turn it upright again. ⑥ Take off the syringe from the bottle ⑦ Place the syringe into your mouth between tongue and cheek, then push the piston slowly until the whole dose has been administered. If the dose to be administered is superior to 11ml, renew operations 2 to 7. . ⑧ Wash the syringe into water after each administration, without dismantling it (do not put it into the dishwasher).</p>

To be filled with the help of healthcare professional**

Informed by

.....

Following

prescription dated

...../...../.....

You can help yourself with **your tracking book* and a calendar** to spot the takes. Never stop or change the rhythm of administration of your treatment without your doctor's advice.

** This file is not a prescription ; refer to your latest prescription..

WHAT TO DO IN CASE OF MISSING OR VOMITING?



- **In case of missing**, do not take the missed dose. The next dose will be taken at the usual time, without doubling it. Please note it in your tracking book*.
- **In case of vomiting**, do not take a new dose. The next dose will be taken at the usual time, without doubling it. Please note it in your tracking book*.

INTERACTIONS WITH OTHERS MEDICINES AND/OR FOOD

Check with your **doctor and your pharmacist** that the medicines (with or without **prescription**) and/or **other substances (food, plants, among which grapefruit and St John's Wort, probiotics, essential oils, food supplements...)** that you are using are compatible with your treatment. Some interactions may have an influence on the efficiency of your treatment or the outcome of side effects.









- You and your partner must use efficient contraceptive methods (condoms and an other method) during the whole treatment.
- Stop breast feeding during the treatment.

*Tracking book available on the sites www.omedibretagne.fr or www.omedit-paysdelaloire.fr

WHAT SIDE EFFECTS MAY OCCUR WITH SPRYCEL® DASATINIB ?

Like all medicines, Dasatinib may bring about side effects, although you may not experience any.

	Side effects	Prevention
	Nauseas, vomitings, loss of appetite	Eat slowly and and make several small light meals. Avoid greasy, fried and spicy food. Rather drink between meals.
	Diarrheas, abdominal pains	Remember to hydrate well : at least 2 litres of water a day. Favor food that is poor in fibers (starchy food, carrots, bananas...). Avoid food that may speed up your intestinal transit (coffee, frozen drinks, dairy, alcohol, raw fruit and vegetables, cereals, fat food, spices...). Rather drink between meals. Watch your weight regularly.
	Swelling of face and limbs	Avoid tight clothes. Keep eating low-salted food. In case of swelling of the legs, raise your feet when sitting and wear support socks or stockings. Watch your weight regularly.
	Unusual bleedings	Use a soft toothbrush to avoid mouth bleeding. Avoid using a rectal thermometer . Do not take aspirin or anti-inflammatory (eg ibuprofen) without medical advice.
	Signs of infection (fever, cough, shivers, sore throat, urinary burns)	Limit contacts with people with a contagious infection. Wash your hands frequently. Disinfect carefully any wound. In case of a fever over 38°C, inform your doctor.
	Pains in muscles	Remember to hydrate frequently. Take moments of rest and light exercise (walking). Avoid repetitive gestures and strenuous effort. In case of important pains, contact your doctor to set up a symptomatic treatment.

Only some of the most frequent side effects are mentionned above. For further information, refer to your packaging or ask your doctor or pharmacist for advice.



You can also ask your questions to your center for side effects monitoring, or notify any side effect on the following site : signalement-sante.gouv.fr



In all cases, remember to make the biological tests prescribed by your doctor.

WHAT TO DO IN CASE OF SIDE EFFECT(S)?

- Take the medication prescribed for side effects in accordance with the doses.
- Respect measures of prevention.

Useful contacts :

Promptly contact your doctor in case of :

- Difficulty to breathe, pains in the chest, sweating
- Bruises, unusual bleedings, blood in vomitings, motions or urines.
- Fever, important shivers.
- Fatigue, confusion, fainting
- Any persisting or worsening side effect.



Remarks :
